

Treating Asymptomatic Bacteriuria: All harm, No Benefit

High Prevalence of Asymptomatic Bacteriuria

- > The bladder is normally colonized in many elderly people
- > A positive urinalysis or culture in the absence of symptoms reveals **colonization, not infection**
- > Treatment of asymptomatic bacteriuria is **not recommended**




It's Hard to Ignore A Positive Test

Habitual Testing + Prevalent Colonization = Unnecessary prescriptions & missing the real diagnosis



Unnecessary Rx and Missed Diagnoses Harm Patients

- > Drug-drug interactions
- > Renal & other complications
- > Increase of multi-drug resistant bacteria
- > *C. difficile* infection
- > Nausea and vomiting
- > Drug allergies
- > Missing the real diagnosis



Myth

Positive urine culture and abnormal urinalysis (positive nitrates or leukocytes, increased white blood cells or pyuria) always indicates a urinary tract infection and requires antibiotics.

Positive urine culture in resident with chronic indwelling catheter always indicates a urinary tract infection and requires antibiotics.

Elderly residents often have a urinary tract infection with no symptoms except a change in mental status or delirium.

In an elderly population, urinary tract infections often present with nonspecific symptoms (e.g., falls, functional decline).

Cloudy or malodorous urine is always diagnostic of a urinary tract infection.

Fact

• Positive urine culture and abnormal urinalysis in a resident without symptoms is consistent with asymptomatic bacteriuria – that is, colonization – not infection. Treatment with antibiotics is not indicated.

• A chronic indwelling catheter is associated with bacteriuria 100% of the time. There is no need to treat unless the resident has symptoms of a UTI.


• A change in mental status or delirium is a non-specific symptom and may accompany a change in condition such as dehydration, constipation, adverse drug effect, pneumonia, urinary retention, metabolic problems, head trauma, environmental changes, or sensory deprivation.
• Mental status change requires an exploration of alternative causes and may not require antibiotics for UTI unless there are more specific signs or symptoms that point to that diagnosis.

Nonspecific symptoms can be seen in many conditions such as dehydration or adverse drug effect. Diagnosing and treating UTIs based on these nonlocalizing symptoms not only results in inappropriate antibiotic use; it also completely misses the real diagnosis.

These changes may be seen in asymptomatic bacteriuria. Other causes can include dehydration, certain medications and diet.

Dangers of Unnecessary Antibiotics

Using Antibiotics Breeds Resistance in Your Patients and the Community



(JID 1999; 179:163)

Infection with Resistant Bacteria Increases Risk of Death

Antibiotic-resistant vs. non-resistant *Klebsiella* bacteria



(ICHE 2008;29:1009-1106)

Do Not Test, Do Not Treat Asymptomatic Bacteriuria¹


Criteria for Urine Testing

Resident without indwelling catheter

- Acute dysuria alone OR
- Fever + at least one of the symptoms below (new or increased) OR
- If no fever, at least two of the symptoms below (new or increased)
 - Gross hematuria
 - Urinary incontinence
 - Urgency
 - Suprapubic pain
 - Costovertebral angle tenderness
 - Frequency

Resident with indwelling catheter

- At least one of the symptoms below (new or increased)
 - Fever
 - Pelvic discomfort
 - Flank pain (back, side pain)
 - Malaise or lethargy no other cause
 - Costovertebral angle (CVA) tenderness
 - Rigors (shaking chills)
 - Delirium
 - Acute hematuria



No symptoms of UTI

- > Do not test urine
- > Do not treat if a urine test was done by someone else or for “routine”

Weakness, delirium, or fever without a focus

- > Individualize care
- > Be mindful of the prevalence of asymptomatic bacteriuria
- > Seek other causes

Specific UTI symptoms

- > Test or treat as usual

Challenges	Strategies for practice change
The resident’s family wants a urine test and antibiotic treatment in the setting of asymptomatic bacteriuria.	<ul style="list-style-type: none"> • Educate the family about the prevalence of asymptomatic bacteriuria, and tell them you do not suspect UTI on clinical grounds. • Emphasize the dangers of antibiotic overuse.
We’ve always ordered urine cultures for nonspecific problems in residents with dementia.	<ul style="list-style-type: none"> • There are many potential causes for nonspecific changes in status and thorough evaluation is needed. • Residents in long-term care frequently have positive urine cultures, even when they are well.
It is okay to give an antibiotic even if it may not be needed. Better safe than sorry.	<ul style="list-style-type: none"> • Antibiotics can cause adverse drug reactions, <i>C. difficile</i> infection, and promote the emergence of multi-drug resistant organisms. They should not be administered unless clinically indicated.
It is hard to ignore a positive urine test even when done for no clearly apparent reason.	<ul style="list-style-type: none"> • Treatment decisions should not be made based on test results alone. • Evaluate the resident clinically and consider a period of observation.

References: ¹ CID 2010;50:625-663; CID 2009;48:149-171; ICHE 2001;22:120-124 CID 2005;40:643-54

Massachusetts Infection Prevention Partnership

Massachusetts Coalition for the Prevention of Medical Errors, Massachusetts Department of Public Health, Massachusetts Senior Care Association, Masspro

Clinical Advisors

Ruth Kandel MD, Director Infection Control, Hebrew Senior Life

Daniel Pallin MD, MPH, Director of Research Brigham & Women’s Hospital Department of Emergency Medicine, and Chairman, Brigham and Women’s Hospital Clinical Investigation Committee

Shira Doron MD, Antimicrobial Steward & Associate Hospital Epidemiologist, Tufts Medical Center

Questions or Copies

froberts@macoalition.org